

## Durham Research Online

---

### Deposited in DRO:

12 December 2019

### Version of attached file:

Accepted Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

The Imbie Team, (2020) 'Mass balance of the Greenland Ice Sheet from 1992 to 2018.', *Nature.*, 579 . pp. 233-239.

### Further information on publisher's website:

<https://doi.org/10.1038/s41586-019-1855-2>

### Publisher's copyright statement:

### Additional information:

---

### Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

# Mass balance of the Greenland Ice Sheet, 1992-2018

The IMBIE Team\*

## Abstract

In recent decades the Greenland Ice Sheet has been a major contributor to global sea-level rise<sup>1,2</sup>, and it is expected to be so in the future<sup>3</sup>. Increases in glacier flow<sup>4-6</sup> and melting from the ice sheet surface<sup>7-9</sup> have been driven by oceanic<sup>10-13</sup> and atmospheric<sup>14,15</sup> warming. Here we compare and combine 26 individual satellite measurements of changes in the ice sheet's volume, flow and gravitational potential to produce a reconciled estimate of its mass balance. Although the ice sheet was close to a state of balance in the 1990's, annual losses rose steadily to peak at  $336 \pm 72$  billion tonnes per year in 2012. In all, Greenland lost  $3928 \pm 341$  billion tonnes of ice between 1992 and 2018, corresponding to an increase in mean sea level of  $10.8 \pm 0.9$  millimetres. Using three regional climate models, we show that reduced surface mass balance has driven  $2028 \pm 509$  billion tonnes (53 %) of the ice loss, owing to increased meltwater runoff.  $1907 \pm 547$  billion tonnes of all ice losses (47 %) are due to increased glacier discharge, which rose from  $30 \pm 34$  billion tonnes per year in the 1990's to  $88 \pm 37$  billion tonnes per year since then. Between 2013 and 2017, the total rate of ice loss slowed to  $217 \pm 19$  billion tonnes per year, on average, as atmospheric circulation favoured cooler conditions<sup>16</sup> and as ocean temperatures fell at the terminus of Jakobshavn Isbræ<sup>17</sup>. Cumulative ice losses from Greenland as a whole have been close to the IPCC's predicted rates for their high-end climate warming scenario<sup>18</sup>, which forecast an additional 70 to 130 millimetres of global sea-level rise by 2100 when compared to their central estimate.

## Introduction

The Greenland Ice Sheet holds enough water to raise mean global sea level by 7.4 m<sup>19</sup>. Its ice flows to the oceans through a network of glaciers and ice streams<sup>20</sup>, each with a substantial inland catchment<sup>21</sup>. Fluctuations in the mass of the Greenland Ice Sheet occur due to variations in snow accumulation, meltwater runoff, ocean-driven melting, and iceberg calving. In recent decades, there have been marked increases in air<sup>22</sup> and ocean<sup>12</sup> temperatures and reductions in summer cloud cover<sup>23</sup> around Greenland. These changes have produced increases in surface runoff<sup>8,24</sup>, supraglacial lake formation<sup>25</sup> and drainage<sup>26</sup>, iceberg calving<sup>27,28</sup>, glacier terminus retreat<sup>29,30</sup>, submarine melting<sup>10,11</sup>, and ice flow<sup>4</sup>, leading to widespread changes in the ice sheet surface elevation, particularly near its margin (Figure 1).

Over recent decades, ice losses from Greenland have made a significant contribution to global sea-level rise<sup>2</sup>, and model projections suggest that this imbalance will continue in a warming climate<sup>3</sup>. Since the early 1990's there have been comprehensive satellite observations of changing ice sheet velocity<sup>4,5,31</sup>, elevation<sup>32-36</sup> and, between 2002 and 2016, its changing gravitational attraction<sup>37,38</sup>, from which complete estimates of Greenland Ice Sheet mass balance are determined<sup>1</sup>. Prior to the 1990's, only partial surveys of the ice sheet elevation<sup>39</sup> and velocity<sup>40</sup> change are available. In combination with models of surface mass balance (the net difference between precipitation, sublimation and meltwater runoff) and glacial isostatic adjustment<sup>41</sup>, satellite measurements have shown a fivefold increase in the rate of ice loss from Greenland overall, rising from  $51 \pm 65$  Gt/yr in the early 1990's to  $263 \pm 30$  Gt/yr between 2005 and 2010<sup>1</sup>. This ice loss has been driven by changes in surface mass balance<sup>7,22</sup> and ice dynamics<sup>6,40</sup>. There was, however, a marked reduction in ice loss between 2013 and 2018, as a consequence of cooler atmospheric conditions and increased precipitation<sup>16</sup>. While the broad pattern of change across Greenland (Figure 1) is one of ice loss,

there is considerable variability; for example, during the 2000's just 4 glaciers were responsible for half of the total ice loss due to increased discharge<sup>6</sup>, whereas many others contribute today<sup>40</sup>. Moreover, some neighbouring ice streams have been observed to speed up over this period while others slowed down<sup>42-44</sup>, suggesting diverse reasons for the changes that have taken place - including their geometrical configuration and basal conditions, as well as the forcing they have experienced<sup>45</sup>. In this study we combine satellite altimetry, gravimetry, and ice velocity measurements to produce a reconciled estimate of the Greenland Ice Sheet mass balance between 1992 and 2018, we evaluate the impact of changes in surface mass balance and uncertainty in glacial isostatic adjustment, and we partition the ice sheet mass loss into signals associated with surface mass balance and ice dynamics. In doing so, we extend a previous assessment<sup>1</sup> to include more satellite and ancillary data and to cover the period since 2012.

## Data and Methods

We use 26 estimates of ice sheet mass balance derived from satellite altimetry (9 data sets), satellite gravimetry (14 data sets) and the input-output method (3 data sets) to assess changes in Greenland ice sheet mass balance. The satellite data were computed using common spatial<sup>21,46</sup> and temporal domains, and using a range of models to estimate signals associated with changes in surface mass balance and glacial isostatic adjustment. Satellite altimetry provides direct measurements of changing ice sheet surface elevation recorded at orbit crossing points<sup>39</sup>, along repeated ground tracks<sup>33</sup>, or using plane-fit solutions<sup>35</sup>, and the ice sheet mass balance is estimated from these measurements either by prescribing the density of the elevation fluctuation<sup>47</sup> or by making an explicit model-based correction for changes in firn height<sup>48</sup>. Satellite gravimetry measures fluctuations in the Earth's gravitational field as computed using either global spherical harmonic solutions<sup>37</sup> or using spatially-discrete mass concentration units<sup>38</sup>. Ice sheet mass changes are determined after making model-based corrections for glacial isostatic adjustment<sup>37</sup>. The input-output method uses model estimates of surface mass balance<sup>7</sup>, which comprises the input, and satellite observations of ice sheet velocity computed from radar<sup>4</sup> and optical<sup>49</sup> imagery combined with airborne measurements of ice thickness<sup>50</sup> to compute changes in marine-terminating glacier discharge into the oceans, which comprises the output. The overall mass balance is the difference between input and output. Not all annual surveys of ice sheet discharge are complete, and sometimes regional extrapolations have to be employed to account for gaps in coverage<sup>40</sup>. Because they provide important ancillary data, we also assess 6 models of glacial isostatic adjustment and 10 models of surface mass balance.

To compare and aggregate the individual satellite data sets, we first adopt a common approach to derive linear rates of ice sheet mass balance over 36-month intervals (see Methods). We then compute error-weighted averages of all altimetry, gravimetry, and input-output group mass trends, and we combine these into a single reconciled estimate of the ice sheet mass balance using error-weighting of the group trends. Uncertainties in individual rates of mass change are estimated as the root sum square of the linear model misfit and their measurement error, uncertainties in group rates are estimated as the root mean square of the contributing time-series errors, and uncertainties in reconciled rates are estimated as their root mean square error divided by the square root of the number of independent groups. Cumulative uncertainties are computed as the root sum square of annual errors, an approach that has been employed in numerous studies<sup>1,18,40,51</sup> and assumes that annual errors are not correlated over time. To improve on this assumption, it will be necessary to consider the covariance of the systematic and random errors present within each mass balance solution (see Methods).

## Inter-comparison of satellite and model results

The satellite gravimetry and satellite altimetry data used in our assessment are corrected for the effects of glacial isostatic adjustment, although the correction is relatively small for altimetry as it appears as a change in elevation and not mass. The most prominent and consistent local signals of glacial isostatic adjustment among the 6 models we have considered are two instances of uplift peaking at about 5-6 mm/yr, one centered over northwest Greenland and Ellesmere Island, and one over northeast Greenland (see Methods and Extended Data Figure 3). Although some models identify a 2 mm/yr subsidence under large parts of the central and southern parts of the ice sheet, it is absent or of lower magnitude in others, which suggests it is less certain (Extended Data Table 1). The greatest difference among model solutions is at Kangerlussuaq Glacier in the southeast where a study<sup>52</sup> has shown that models and observations agree if a localized weak Earth structure associated with overpassing the Iceland hotspot is assumed; the effect is to offset earlier estimates of mass trends associated with glacial isostatic adjustment by about 20 Gt/yr. Farther afield, the highest spread between modelled uplift occurs on Baffin Island and beyond due to variations in regional model predictions related to the demise of the Laurentide Ice Sheet<sup>52,53</sup>. This regional uncertainty is likely a major factor in the spread across the ice-sheet-wide estimates. Nevertheless, at  $-3 \pm 20$  Gt/yr, the mass signal associated with glacial isostatic adjustment in Greenland shows no coherent substantive change and is negligible relative to reported ice sheet mass trends<sup>1</sup>.

There is generally good agreement between the models of Greenland Ice Sheet surface mass balance that we have assessed for determining mass input - particularly those of a similar class; for example, 70% of all model estimated of runoff and accumulation fall within 1-sigma of their mean (see Methods and Extended Data Table 2). The exceptions are a global reanalysis with coarse spatial resolution that tends to underestimate runoff due to its poor delineation of the ablation zone, and a snow process model that tends to underestimate precipitation and to overestimate runoff in most sectors. Among the other 8 models, the average surface mass balance between 1980 and 2012 is  $361 \pm 40$  Gt/yr, with a marked negative trend over time (Extended Data Figure 4) mainly due to increased runoff<sup>7</sup>. At regional scale, the largest differences occur in the northeast, where two regional climate models predict significantly less runoff, and in the southeast, where there is considerable spread in precipitation and runoff across all models. All models show high temporal variability in surface mass balance components, and all models show that the southeast receives the highest net intake of mass at the surface due to high rates of snowfall originating from the Icelandic Low<sup>54</sup>. By contrast, the southwest, which features the widest ablation zone<sup>7</sup>, has experienced alternate periods of net surface mass loss and gain over recent decades, and has the lowest average surface mass balance across the ice sheet.

We assessed the consistency of the satellite altimetry, gravimetry, and input-output method estimates of Greenland Ice Sheet mass balance using common spatial and temporal domains (see Figure 2 and Methods). In general, there is close agreement between estimates determined using each approach, and the standard deviations of coincident altimetry, gravimetry, and input-output method annual mass balance solutions are 31, 26, and 22 Gt/yr, respectively (Extended Data Table 3). Once averages were formed for each technique, the resulting estimates of mass balance were also closely aligned (e.g. Extended Data Figure 6). For example, over the common period 2005 to 2015, the average Greenland Ice Sheet mass balance is  $-249 \pm 61$  Gt/yr and, by comparison, the spread of the altimetry, gravimetry, and input-output method estimates is just 36 Gt/yr (Extended Data Table 3). The estimated uncertainty of the aggregated mass balance solution (see Methods) is larger than the standard deviation of model corrections for glacial isostatic adjustment (20 Gt/yr for gravimetry) and for surface mass balance (40 Gt/yr), which suggests that their collective impacts

have been adequately compensated, and it is also larger than the estimated 30 Gt/yr mass losses from peripheral ice caps<sup>55,56</sup>, which are not accounted for in all individual solutions. In keeping with results from Antarctica<sup>51</sup>, rates of mass loss determined using the input-output method are the most negative, and those determined from altimetry are the least negative. However, the spread among the three techniques is 5 times lower for Greenland than it is for Antarctica<sup>51</sup>, reflecting differences in the ice sheet size, the complexity of the mass balance processes, and limitations of the various geodetic techniques.

## Ice sheet mass balance

We aggregated the average mass balance estimates from gravimetry, altimetry and the input-output method to form a single, time-varying record (Figure 2) and then integrated these data to determine the cumulative mass lost from Greenland since 1992 (Figure 3). Although Greenland has been losing ice throughout most of the intervening period, the rate of loss has varied significantly. Between 1992 and 2012, the rate of ice loss progressively increased, reaching a maximum of  $336 \pm 72$  Gt/yr in 2012, coinciding with the extreme summertime surface melting that occurred in that year<sup>57</sup>. Since 2012, however, the trend has reversed, with a progressive reduction in the rate of mass loss during the subsequent period. By 2018 – the last complete year of our survey – the annual rate of ice mass loss had reduced to  $144 \pm 57$  Gt/yr. The highly variable nature of ice losses from Greenland is a consequence of the wide range of physical processes that are affecting different sectors of the ice sheet<sup>17,35,44</sup>, which suggests that care should be taken when extrapolating sparse measurements in space or time. Although the rates of mass loss we have computed between 1992 and 2011 are 18 % less negative than those of a previous assessment, which included far fewer data sets<sup>1</sup>, the results are consistent given their respective uncertainties. Altogether, the Greenland Ice Sheet has lost  $3928 \pm 341$  Gt of ice to the ocean since 1992, with roughly half of this loss occurring during the 6-year period between 2006 and 2012.

To determine the proportion of mass lost due to surface and ice dynamical processes, we computed the contemporaneous trend in Greenland Ice Sheet surface mass balance - the net balance between precipitation and ablation<sup>7</sup>, which is controlled by interactions with the atmosphere (Figure 3). In Greenland, recent trends in surface mass balance have been largely driven by meltwater runoff<sup>54</sup>, which has increased as the regional climate has warmed<sup>14</sup>. Because direct observations of ice sheet surface mass balance are too scarce to provide full temporal and spatial coverage<sup>58</sup>, regional estimates are usually taken from atmospheric models that are evaluated with existing observations. Our evaluation (see Methods) shows that the finer spatial resolution regional climate models produce consistent results, likely due to their ability to capture local changes in melting and precipitation associated with atmospheric forcing, and to resolve the full extent of the ablation zone<sup>59</sup>. We therefore compare and combine estimates of Greenland surface mass balance derived from three regional climate models; RACMO2.3p2<sup>59</sup>, MARv3.6<sup>22</sup> and HIRHAM<sup>9</sup>. To assess the surface mass change across the Greenland Ice Sheet between 1980 and 2018, we accumulate surface mass balance anomalies from each of the regional climate models (Extended Data Figure 7) and average them into a single estimate (Figure 3). Surface mass balance anomalies are computed with respect to the average between 1980 and 1990, which corresponds to a period of approximate balance<sup>8</sup> and is common to all models. In this comparison, all three models show that the Greenland Ice Sheet entered abruptly into a period of anomalously low surface mass balance in the late 1990's and, when combined, they show that the ice sheet lost  $2028 \pm 509$  Gt of its mass due to meteorological processes between 1992 and 2018 (Table 1).

Just over half (53 %) of all mass losses from Greenland – and much of their short-term variability – have been due to variations in the ice sheet’s surface mass balance and its indirect impacts on firn processes. For example, between 2007 and 2012, 72 % of the total ice loss ( $193 \pm 30$  Gt/yr) was due to surface mass balance, compared to 36 % ( $87 \pm 44$  Gt/yr) over the preceding 15 years and 58 % ( $139 \pm 25$  Gt/yr) since then (Table 1). The rise in the total rate of ice loss during the late-2000s coincided with warmer atmospheric conditions, which promoted several episodes of widespread melting and runoff<sup>15,60</sup>. The reduction in surface mass loss since then is associated with a shift of the North Atlantic Oscillation, which brought about cooler atmospheric conditions and increased precipitation along the southeastern coast<sup>16</sup>. Trends in the total ice sheet mass balance are not, however, entirely due to surface mass balance and, by differencing these two signals, we can estimate the total change in mass loss due to ice dynamical imbalance – i.e. the integrated, net mass loss from those glaciers whose velocity does not equal their long-term mean (Figure 3). Although this approach is indirect, it makes use of all the satellite observations and regional climate models included in our study, overcoming limitations in the spatial and temporal sampling of ice discharge estimates derived from ice velocity and thickness data. Our estimate shows that, between 1992 and 2018, Greenland lost  $1907 \pm 547$  Gt of ice due to the dynamical imbalance of glaciers relative to their steady state, accounting for 47 % of the total imbalance (Table 1). Losses due to increased ice discharge rose sharply in the early 2000’s when Jakobshavn Isbræ<sup>61-63</sup> and several other outlet glaciers in the southeast<sup>64-66</sup> sped up, and the discharge losses are now four times higher than in the 1990’s. For a period between 2002 and 2007, ice dynamical imbalance was the major source of ice loss from the ice sheet as a whole, although the situation has since returned to be dominated by surface mass losses as several glaciers have slowed down<sup>17,30</sup>.

Despite a reduction in the overall rate of ice loss from Greenland between 2013 and 2018 (Figure 2), the ice sheet mass balance remained negative, adding  $10.8 \pm 0.9$  mm to global sea level since 1992. Although the average sea level contribution is  $0.42 \pm 0.08$  mm/yr, the five-year average rate varied by a factor 5 over the 25-year period, peaking at  $0.75 \pm 0.05$  mm/yr between 2007 and 2012. The variability in Greenland ice loss illustrates the importance of accounting for yearly fluctuations when attempting to close the global sea level budget<sup>2,67</sup>. Satellite records of ice sheet mass balance are also an important tool for evaluating numerical models of ice sheet evolution<sup>68</sup>. In their 2013 assessment, the Intergovernmental Panel on Climate Change (IPCC) predicted ice losses from Greenland due to surface mass balance and glacier dynamics under a range of scenarios, beginning in 2007<sup>18</sup> (Figure 4). Although ice losses from Greenland have fluctuated considerably during the 12-year period of overlap between the IPCC predictions and our reconciled time series, the total change and average rate ( $0.70$  mm/yr) are close to the upper range predictions (0.74 mm/yr), which implies a 70 to 130 mm of sea-level rise by the year 2100 above central estimates. The drop in ice losses between 2013 and 2018, however, shifted rates towards the lower end projections, and a longer period of comparison is required to establish whether the upper trajectory will continue to be followed. Even greater sea level contribution cannot be ruled out if feedbacks between the ice sheet and other elements of the climate system are underestimated by current ice sheet models<sup>3</sup>. Although the volume of ice stored in Greenland is a small fraction of that in Antarctica (12 %), its recent losses have been ~38 % higher<sup>51</sup> as a consequence of the relatively strong atmospheric<sup>14,15</sup> and oceanic<sup>10,11</sup> warming that has occurred in its vicinity, and its status as a major source of sea-level rise is expected to continue<sup>3,18</sup>.

## Conclusions

We combine 26 satellite estimates of ice sheet mass balance and assess 10 models of ice sheet surface mass balance and 6 models of glacial isostatic adjustment, to show that the Greenland Ice



Sheet lost  $3928 \pm 341$  Gt of ice between 1992 and 2018. During the common period 2005 to 2015, the spread of mass balance estimates derived from satellite altimetry, gravimetry, and the input-output method is  $32$  Gt/yr, or  $13\%$  of the estimated rate of imbalance. The rate of ice loss has generally increased over time, rising from  $20 \pm 24$  Gt/yr between 1992 to 1997, peaking at  $269 \pm 17$  Gt/yr between 2007 and 2012, and reducing to  $238 \pm 20$  Gt/yr between 2012 and 2017. Just over half ( $2028 \pm 509$  Gt, or  $53\%$ ) of the ice losses are due to reduced surface mass balance (mostly meltwater runoff) associated with changing atmospheric conditions<sup>14,15,69</sup>, and these changes have also driven the shorter-term temporal variability in ice sheet mass balance. Despite variations in the imbalance of individual glaciers<sup>5,6,40</sup>, ice losses due to increasing discharge from the ice sheet as a whole have risen steadily from  $26 \pm 35$  Gt/yr in the 1990's to  $88 \pm 37$  Gt/yr since then, and account for just under half of all losses (47 %) over the survey period.

Our assessment shows that estimates of Greenland Ice Sheet mass balance derived from satellite altimetry, gravimetry, and the input-output method agree to within 20 Gt/yr, that model estimates of surface mass balance agree to within 40 Gt/yr, and that model estimates of glacial isostatic adjustment agree to within 20 Gt/yr. These differences represent a small fraction ( $13\%$ ) of the Greenland Ice Sheet mass imbalance and are comparable to its estimated uncertainty ( $28$  Gt/yr). Nevertheless, there is still departure among models of glacial isostatic adjustment in northern Greenland. Spatial resolution is a key factor in the degree to which models of surface mass balance can represent ablation and precipitation at local scales, and estimates of ice sheet mass balance determined from satellite altimetry and the input-output method continue to be positively and negatively biased, respectively, compared to those based on satellite gravimetry (albeit by small amounts). More satellite estimates of ice sheet mass balance at the start (1990's) and end (2010's) of our record would help to reduce the dependence on fewer data during those periods; although new missions<sup>70,71</sup> will no doubt address the latter, further analysis of historical satellite data is required to address the former.

## Acknowledgements

This work is an outcome of the Ice Sheet Mass Balance Inter-Comparison Exercise (IMBIE) supported by the ESA Climate Change Initiative and the NASA Cryosphere Program. A.S. was additionally supported by a Royal Society Wolfson Research Merit Award.

## Author Contributions

A.S. and E.I. designed and led the study. E.R., B.S., M.v.d.B., I.V. and P.W. led the input-output method, altimetry, surface mass balance (SMB), gravimetry and glacial isostatic adjustment (GIA) experiments, respectively. A.S., E.I., K.B., M.E., A.H., I.J., G.K., S.N., T.P., E.R., T.Sc., N.S., B.S., M.v.d.B., I.V., T.W., and P.W. supervised the assessment exercise. G.M., M.E.P., and T.Sl. performed the mass balance data collation and analysis. T.Sl. performed the AR5 data analysis. P.W. and I.S. performed the GIA data analysis. M.v.W. and T.Sl. performed the SMB data analysis. A.S., E.I., K.B., M.E., N.G., A.H., H.K., M.M., I.O., I.S., T.Sl., M.v.W., and P.W. wrote the manuscript. A.S., K.B., H.K., G.M., M.E.P., I.S., S.B.S., T.Sl., P.W., and M.v.W. prepared the figures and tables. All authors participated in the data interpretation and commented on the manuscript.

## Competing Interests

The authors declare no competing interests.

## 266 The IMBIE Team

267 Andrew Shepherd<sup>1\*</sup>, Erik Ivins<sup>2</sup>, Eric Rignot<sup>2,3</sup>, Ben Smith<sup>4</sup>, Michiel van den Broeke<sup>5</sup>, Isabella  
 268 Velicogna<sup>2,3</sup>, Pippa Whitehouse<sup>6</sup>, Kate Briggs<sup>1</sup>, Ian Joughin<sup>4</sup>, Gerhard Krinner<sup>7</sup>, Sophie Nowicki<sup>8</sup>, Tony  
 269 Payne<sup>9</sup>, Ted Scambos<sup>10</sup>, Nicole Schlegel<sup>2</sup>, Geruo A<sup>3</sup>, Cécile Agosta<sup>11</sup>, Andreas Ahlstrøm<sup>12</sup>, Greg  
 270 Babonis<sup>13</sup>, Valentina R. Barletta<sup>14</sup>, Anders A. Bjørk<sup>15</sup>, Alejandro Blazquez<sup>16</sup>, Jennifer Bonin<sup>17</sup>, William  
 271 Colgan<sup>12</sup>, Beata Csatho<sup>13</sup>, Richard Cullather<sup>18</sup>, Marcus Engdahl<sup>19</sup>, Denis Felikson<sup>8</sup>, Xavier Fettweis<sup>11</sup>,  
 272 Rene Forsberg<sup>14</sup>, Anna E. Hogg<sup>1</sup>, Hubert Gallee<sup>7</sup>, Alex Gardner<sup>2</sup>, Lin Gilbert<sup>20</sup>, Noel Gourmelen<sup>21</sup>,  
 273 Andreas Groh<sup>22</sup>, Brian Gunter<sup>23</sup>, Edward Hanna<sup>24</sup>, Christopher Harig<sup>25</sup>, Veit Helm<sup>26</sup>, Alexander  
 274 Horvath<sup>27</sup>, Martin Horwath<sup>22</sup>, Shfaqat Khan<sup>14</sup>, Kristian K. Kjeldsen<sup>12,28</sup>, Hannes Konrad<sup>29</sup>, Peter L.  
 275 Langen<sup>30</sup>, Benoit Lecavalier<sup>31</sup>, Bryant Loomis<sup>8</sup>, Scott Luthcke<sup>8</sup>, Malcolm McMillan<sup>32</sup>, Daniele Melini<sup>33</sup>,  
 276 Sebastian Mernild<sup>34,35,36,37</sup>, Yara Mohajerani<sup>3</sup>, Philip Moore<sup>38</sup>, Ruth Mottram<sup>30</sup>, Jeremie Mouginot<sup>3,7</sup>,  
 277 Gorka Moyano<sup>39</sup>, Alan Muir<sup>20</sup>, Thomas Nagler<sup>40</sup>, Grace Nield<sup>6</sup>, Johan Nilsson<sup>2</sup>, Brice Noël<sup>5</sup>, Ines  
 278 Ootosaka<sup>1</sup>, Mark E. Pattle<sup>39</sup>, W. Richard Peltier<sup>41</sup>, Nadège Pie<sup>42</sup>, Roelof Rietbroek<sup>43</sup>, Helmut Rott<sup>40</sup>,  
 279 Louise Sandberg Sørensen<sup>14</sup>, Ingo Sasgen<sup>26</sup>, Himanshu Save<sup>42</sup>, Bernd Scheuchl<sup>3</sup>, Ernst Schrama<sup>44</sup>,  
 280 Ludwig Schröder<sup>22,26</sup>, Ki-Weon Seo<sup>45</sup>, Sebastian B. Simonsen<sup>14</sup>, Thomas Slater<sup>1</sup>, Giorgio Spada<sup>46</sup>, Tyler  
 281 Sutterley<sup>3</sup>, Matthieu Talpe<sup>2</sup>, Lev Tarasov<sup>31</sup>, Willem Jan van de Berg<sup>5</sup>, Wouter van der Wal<sup>47</sup>, Melchior  
 282 van Wessem<sup>5</sup>, Bramha Dutt Vishwakarma<sup>48</sup>, David Wiese<sup>2</sup>, David Wilton<sup>49</sup>, Thomas Wagner<sup>50</sup>, Bert  
 283 Wouters<sup>5,47</sup> & Jan Wuite<sup>40</sup>

284


285 <sup>1</sup>Centre for Polar Observation and Modelling, University of Leeds, Leeds, UK. <sup>2</sup>NASA Jet Propulsion  
 286 Laboratory, California Institute of Technology, Pasadena, CA, USA. <sup>3</sup>Department of Earth System  
 287 Science, University of California, Irvine, CA, USA. <sup>4</sup>Department of Earth and Space Sciences,  
 288 University of Washington, Seattle, WA, USA. <sup>5</sup>Institute for Marine and Atmospheric Research,  
 289 Utrecht University, Utrecht, The Netherlands. <sup>6</sup>Department of Geography, Durham University,  
 290 Durham, UK. <sup>7</sup>Institute of Environmental Geosciences, Université Grenoble Alpes, Grenoble, France.  
 291 <sup>8</sup>Cryospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, USA. <sup>9</sup>School  
 292 of Geographical Sciences, University of Bristol, Bristol, UK. <sup>10</sup>Earth Science and Observation Center,  
 293 University of Colorado, Boulder, CO, USA. <sup>11</sup>Department of Geography, University of Liège, Liège,  
 294 Belgium. <sup>12</sup>Geological Survey of Denmark and Greenland, Copenhagen, Denmark. <sup>13</sup>Department of  
 295 Geology, State University of New York at Buffalo, Buffalo, NY, USA. <sup>14</sup>DTU Space, National Space  
 296 Institute, Technical University of Denmark, Kongens Lyngby, Denmark. <sup>15</sup>Department of Geosciences  
 297 and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark. <sup>16</sup>LEGOS,  
 298 Université de Toulouse, Toulouse, France. <sup>17</sup>College of Marine Sciences, University of South Florida,  
 299 Tampa, FL, USA. <sup>18</sup>Global Modeling and Assimilation Office, NASA Goddard Space Flight Center,  
 300 Greenbelt, MD, USA. <sup>19</sup>ESA-ESRIN, Frascati, Italy. <sup>20</sup>Mullard Space Science Laboratory, University  
 301 College London, Holmbury St Mary, UK. <sup>21</sup>School of Geosciences, University of Edinburgh, Edinburgh,  
 302 UK. <sup>22</sup>Institute for Planetary Geodesy, Technische Universität Dresden, Dresden, Germany. <sup>23</sup>Daniel  
 303 Guggenheim School of Aerospace Engineering, Georgia Institute of Technology, Atlanta, GA, USA.  
 304 <sup>24</sup>School of Geography, University of Lincoln, Lincoln, UK. <sup>25</sup>Department of Geosciences, University of  
 305 Arizona, Tucson, AZ, USA. <sup>26</sup>Alfred Wegener Institute, Helmholtz Centre for Polar and Marine  
 306 Research, Bremerhaven, Germany. <sup>27</sup>Institute of Astronomical and Physical Geodesy, Technical  
 307 University Munich, Munich, Germany. <sup>28</sup>GeoGenetics, Globe Institute, University of Copenhagen,  
 308 Copenhagen, Denmark. <sup>29</sup>Deutscher Wetterdienst, Offenbach, Germany. <sup>30</sup>Danish Meteorological  
 309 Institute, Copenhagen, Denmark. <sup>31</sup>Department of Physics and Physical Oceanography, Memorial  
 310 University of Newfoundland, St. Johns, Newfoundland and Labrador, Canada. <sup>32</sup>University of  
 311 Lancaster, Lancaster, UK. <sup>34</sup>Istituto Nazionale di Geofisica e Vulcanologia, Roma, Italy <sup>34</sup>Nansen  
 312 Environmental and Remote Sensing Centre, Bergen, Norway. <sup>35</sup>Faculty of Engineering and Science,  
 313 Western Norway University of Applied Sciences, Sogndal, Norway. <sup>36</sup>Direction of Antarctic and Sub-  
 314 Antarctic Programs, Universidad de Magallanes, Punta Arenas, Chile, <sup>37</sup>Geophysical Institute,  
 315 University of Bergen, Norway. <sup>38</sup>School of Engineering, Newcastle University, Newcastle upon Tyne,



316 UK. <sup>39</sup>isardSAT, Barcelona, Spain. <sup>40</sup>ENVEO, Innsbruck, Austria. <sup>41</sup>Department of Physics, University of  
317 Toronto, Toronto, Ontario, Canada. <sup>42</sup>Center for Space Research, University of Texas, Austin, TX, USA.  
318 <sup>43</sup>Institute of Geodesy and Geoinformation, University of Bonn, Bonn, Germany. <sup>44</sup>Department of  
319 Space Engineering, Delft University of Technology, Delft, The Netherlands. <sup>45</sup>Department of Earth  
320 Science Education, Seoul National University, Seoul, South Korea. <sup>46</sup>Dipartimento di Scienze Pure e  
321 Applicate, Università di Urbino "Carlo Bo", Italy. <sup>47</sup>Department of Civil Engineering, Delft University of  
322 Technology, Delft, The Netherlands. <sup>48</sup>Geodetic Institute, University of Stuttgart, Stuttgart, Germany.  
323 <sup>49</sup>Department of Computer Science, University of Sheffield, UK. <sup>50</sup>NASA Headquarters, Washington  
324 D.C., USA.


325 \*Corresponding author: Andrew Shepherd [a.shepherd@leeds.ac.uk](mailto:a.shepherd@leeds.ac.uk)

326




328


329



**Figure 2 | Greenland Ice Sheet mass balance.** Rate of mass change ( $dM/dt$ ) of the Greenland Ice Sheet as determined from the satellite-altimetry (red), input-output method (blue) and gravimetry (green) assessments included in this study. In each case,  $dM/dt$  is computed at annual intervals from time series of relative mass change using a three-year window. An average of estimates across each class of measurement technique is also shown for each year (black). The estimated  $1\sigma$ ,  $2\sigma$  and  $3\sigma$  ranges of the class average is shaded in dark, mid and light grey, respectively; 97 % of all estimates fall within the  $1\sigma$  range, given their estimated individual errors. The equivalent sea level contribution of the mass change is also indicated, and the number of individual mass-balance estimates collated at each epoch is shown below each chart entry.



**Figure 3 | Cumulative anomalies in Greenland Ice Sheet total mass, surface mass balance and ice dynamics.** The total change (dark blue) is determined as the integral of the average rate of ice sheet mass change (Figure 2). The change in surface mass balance (green) is determined from three regional climate models relative to their mean over the period 1980-1990. The change associated with ice dynamics (light blue) is determined as the difference between the change in total and surface mass. The estimated 1σ uncertainties of the cumulative changes are shaded. The dotted line shows the result of a previous assessment <sup>1</sup>. The equivalent sea level contribution of the mass change is also indicated. Vertical lines mark consecutive five-year epochs since the start of our satellite record in 1992.



**Figure 4 | Observed and predicted sea level contribution due to Greenland Ice Sheet mass change.** The global sea-level contribution from Greenland Ice Sheet mass change according to this study (black line) and IPCC AR5 projections between 1992–2040 (left) and 2040–2100 (right) including upper (red), mid (orange), and lower (blue) estimates from the sum of modelled surface mass balance and rapid ice dynamical contributions. Darker coloured lines represent pathways from the five AR5 scenarios in order of increasing emissions: RCP2.6, RCP4.5, RCP6.0, SRES A1B and RCP8.5. Shaded areas represent the spread of AR5 emissions scenarios and the 1 $\sigma$  estimated error on the IMBIE data. The bar chart plot (inset) shows the average annual rates of sea-level rise (in mm/yr) during the overlap period 2007–2018 and their standard deviations. Cumulative AR5 projections have been offset to make them equal to the observational record at their start date (2007).

Region	1992-1997 (Gt/yr)	1997-2002 (Gt/yr)	2002-2007 (Gt/yr)	2007-2012 (Gt/yr)	2012-2017 (Gt/yr)	1992-2011 (Gt/yr)	1992-2018 (Gt/yr)
Total	-20 ± 24	-48 ± 25	-175 ± 20	-269 ± 17	-238 ± 20	-117 ± 24	-150 ± 28
Surface	6 ± 22	-15 ± 25	-78 ± 29	-193 ± 30	-139 ± 25	-62 ± 31	-79 ± 33
Dynamics	-26 ± 35	-33 ± 31	-97 ± 34	-76 ± 34	-100 ± 29	-55 ± 38	-70 ± 40

**Table 1 | Rates of Greenland Ice Sheet total, surface, and dynamical mass change.** Total rates were determined from all satellite measurements over various epochs, rates of surface mass change were determined from three regional climate models, and rates of dynamical mass change were determined as the difference. The period 1992–2011 is included for comparison to a previous assessment <sup>1</sup>, which reported a mass-balance estimate of -142 ± 49 Gt/yr based on far fewer data. The small differences in our updated estimate is due to our inclusion of more data and an updated aggregation

scheme (see Methods). Errors are  $1\sigma$ .

## References

- 1 Shepherd, A. et al. A reconciled estimate of ice-sheet mass balance. *Science* 338, 1183-1189, doi:10.1126/science.1228102 (2012).
- 2 Cazenave, A. et al. Global sea-level budget 1993-present. *Earth Sys. Sci. Data* 10, 1551-1590, doi:10.5194/essd-10-1551-2018 (2018).
- 3 Pattyn, F. et al. The Greenland and Antarctic ice sheets under 1.5 °C global warming. *Nat. Clim. Change* 8, 1053-1061, doi:10.1038/s41558-018-0305-8 (2018).
- 4 Rignot, E. & Kanagaratnam, P. Changes in the velocity structure of the Greenland ice sheet. *Science* 311, 986-990 (2006).
- 5 Moon, T., Joughin, I., Smith, B. & Howat, I. 21st-century evolution of Greenland outlet glacier velocities. *Science* 336, 576-578, doi:10.1126/science.1219985 (2012).
- 6 Enderlin, E. M. et al. An improved mass budget for the Greenland ice sheet. *Geophysical Research Letters* 41, 866-872, doi:10.1002/2013GL059010 (2014).
- 7 Van Den Broeke, M. et al. Partitioning recent Greenland mass loss. *Science* 326, 984-986, doi:10.1126/science.1178176 (2009).
- 8 Trusel, L. D. et al. Nonlinear rise in Greenland runoff in response to post-industrial Arctic warming. *Nature* 564, 104-108, doi:10.1038/s41586-018-0752-4 (2018).
- 9 Lucas-Picher, P. et al. Very high resolution regional climate model simulations over Greenland: Identifying added value. *J. Geophys. Res. D Atmos.* 117, doi:10.1029/2011JD016267 (2012).
- 10 Holland, D. M., Thomas, R. H., De Young, B., Ribergaard, M. H. & Lyberth, B. Acceleration of Jakobshavn Isbrae triggered by warm subsurface ocean waters. *Nature Geoscience* 1, 659-664 (2008).
- 11 Seale, A., Christoffersen, P., Mugford, R. I. & O'Leary, M. Ocean forcing of the Greenland Ice Sheet: Calving fronts and patterns of retreat identified by automatic satellite monitoring of eastern outlet glaciers. *J. Geophys. Res. F Earth Surf.* 116, doi:10.1029/2010JF001847 (2011).
- 12 Straneo, F. & Heimbach, P. North Atlantic warming and the retreat of Greenland's outlet glaciers. *Nature* 504, 36-43, doi:10.1038/nature12854 (2013).
- 13 Rignot, E., Fenty, I., Menemenlis, D. & Xu, Y. Spreading of warm ocean waters around Greenland as a possible cause for glacier acceleration. *Annals of Glaciology* 53, 257-266, doi:10.3189/2012AoG60A136 (2012).
- 14 Hanna, E., Mernild, S. H., Cappelen, J. & Steffen, K. Recent warming in Greenland in a long-term instrumental (1881-2012) climatic context: I. Evaluation of surface air temperature records. *Environ. Res. Lett.* 7, doi:10.1088/1748-9326/7/4/045404 (2012).
- 15 Fettweis, X. et al. Brief communication Important role of the mid-tropospheric atmospheric circulation in the recent surface melt increase over the Greenland ice sheet. *Cryosphere* 7, 241-248, doi:10.5194/tc-7-241-2013 (2013).



- 376 16 Bevis, M. et al. Accelerating changes in ice mass within Greenland, and the ice sheet's  
377 sensitivity to atmospheric forcing. *Proceedings of the National Academy of Sciences of the*  
378 *United States of America* 116, 1934-1939, doi:10.1073/pnas.1806562116 (2019).
- 379 17 Khazendar, A. et al. Interruption of two decades of Jakobshavn Isbrae acceleration and  
380 thinning as regional ocean cools. *Nature Geoscience* 12, 277-283, doi:10.1038/s41561-019-  
381 0329-3 (2019).
- 382 18 Church, J. A. et al. in *Climate Change 2013: The Physical Science Basis. Contribution of*  
383 *Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*  
384 *Change* (eds T. F. Stocker et al.) Ch. 13, 1137–1216 (Cambridge University Press, 2013).
- 385 19 Morlighem, M. et al. BedMachine v3: Complete Bed Topography and Ocean Bathymetry  
386 Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation.  
387 *Geophysical Research Letters* 44, 11,051-011,061, doi:10.1002/2017GL074954 (2017).
- 388 20 Joughin, I., Smith, B. E., Howat, I. M., Scambos, T. & Moon, T. Greenland flow variability from  
389 ice-sheet-wide velocity mapping. *Journal of Glaciology* 56, 415-430,  
390 doi:10.3189/002214310792447734 (2010).
- 391 21 Zwally, H. J., Giovinetto, M. B., Beckley, M. A. & Saba, J. L. Antarctic and Greenland drainage  
392 systems. GSFC Cryospheric Sciences Laboratory.  
393 [https://icesat4.gsfc.nasa.gov/cryo\\_data/ant\\_grn\\_drainage\\_systems.php](https://icesat4.gsfc.nasa.gov/cryo_data/ant_grn_drainage_systems.php) (2012).
- 394 22 Fettweis, X. et al. Reconstructions of the 1900-2015 Greenland ice sheet surface mass  
395 balance using the regional climate MAR model. *Cryosphere* 11, 1015-1033, doi:10.5194/tc-  
396 11-1015-2017 (2017).
- 397 23 Hofer, S., Tedstone, A. J., Fettweis, X. & Bamber, J. L. Decreasing cloud cover drives the  
398 recent mass loss on the Greenland Ice Sheet. *Sci. Adv.* 3, doi:10.1126/sciadv.1700584 (2017).
- 399 24 Van Den Broeke, M. R. et al. On the recent contribution of the Greenland ice sheet to sea  
400 level change. *Cryosphere* 10, 1933-1946, doi:10.5194/tc-10-1933-2016 (2016).
- 401 25 Leeson, A. A. et al. Supraglacial lakes on the Greenland ice sheet advance inland under  
402 warming climate. *Nat. Clim. Change* 5, 51-55, doi:10.1038/nclimate2463 (2015).
- 403 26 Palmer, S., McMillan, M. & Morlighem, M. Subglacial lake drainage detected beneath the  
404 Greenland ice sheet. *Nat. Commun.* 6, doi:10.1038/ncomms9408 (2015).
- 405 27 Nick, F. M. et al. The response of Petermann Glacier, Greenland, to large calving events, and  
406 its future stability in the context of atmospheric and oceanic warming. *Journal of Glaciology*  
407 58, 229-239, doi:10.3189/2012JoG11J242 (2012).
- 408 28 Amundson, J. M. et al. Glacier, fjord, and seismic response to recent large calving events,  
409 Jakobshavn Isbræ, Greenland. *Geophysical Research Letters* 35, doi:10.1029/2008GL035281  
410 (2008).
- 411 29 Joughin, I. et al. Ice-front variation and tidewater behavior on Helheim and Kangerdlugssuaq  
412 Glaciers, Greenland. *J. Geophys. Res. F Earth Surf.* 113, doi:10.1029/2007JF000837 (2008).
- 413 30 Lemos, A. et al. Ice velocity of Jakobshavn Isbræ, Petermann Glacier, Nioghalvfjærdsfjorden,  
414 and Zachariæ Isstrøm, 2015-2017, from Sentinel 1-a/b SAR imagery. *Cryosphere* 12, 2087-  
415 2097, doi:10.5194/tc-12-2087-2018 (2018).

416 31 Nagler, T., Rott, H., Hetzenecker, M., Wuite, J. & Potin, P. The Sentinel-1 mission: New  
417 opportunities for ice sheet observations. *Remote Sens.* 7, 9371-9389,  
418 doi:10.3390/rs70709371 (2015).

419 32 Krabill, W. et al. Greenland ice sheet: High-elevation balance and peripheral thinning.  
420 *Science* 289, 428-430 (2000).

421 33 Pritchard, H. D., Arthern, R. J., Vaughan, D. G. & Edwards, L. A. Extensive dynamic thinning  
422 on the margins of the Greenland and Antarctic ice sheets. *Nature* 461, 971-975 (2009).

423 34 Kjeldsen, K. K. et al. Spatial and temporal distribution of mass loss from the Greenland Ice  
424 Sheet since AD 1900. *Nature* 528, 396-400, doi:10.1038/nature16183 (2015).

425 35 McMillan, M. et al. A high-resolution record of Greenland mass balance. *Geophysical*  
426 *Research Letters* 43, 7002-7010, doi:10.1002/2016GL069666 (2016).

427 36 Sandberg Sørensen, L. et al. 25 years of elevation changes of the Greenland Ice Sheet from  
428 ERS, Envisat, and CryoSat-2 radar altimetry. *Earth and Planetary Science Letters* 495, 234-  
429 241, doi:10.1016/j.epsl.2018.05.015 (2018).

430 37 Velicogna, I. & Wahr, J. Greenland mass balance from GRACE. *Geophysical Research Letters*  
431 32, art-L18505 (2005).

432 38 Luthcke, S. B. et al. Recent Greenland Ice Mass Loss by Drainage System from Satellite  
433 Gravity Observations. *Science* 314, 1286-1289 (2006).

434 39 Zwally, H. J., Brenner, A. C., Major, J. A., Bindshadler, R. A. & Marsh, J. G. Growth of  
435 Greenland ice sheet: Measurement. *Science* 246, 1587-1589,  
436 doi:10.1126/science.246.4937.1587 (1989).

437 40 Mouginot, J. et al. Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018.  
438 *Proceedings of the National Academy of Sciences of the United States of America* (2019).

439 41 Lecavalier, B. S. et al. A model of Greenland ice sheet deglaciation constrained by  
440 observations of relative sea level and ice extent. *Quaternary Science Reviews* 102, 54-84,  
441 doi:10.1016/j.quascirev.2014.07.018 (2014).

442 42 Felikson, D. et al. Inland thinning on the Greenland ice sheet controlled by outlet glacier  
443 geometry. *Nature Geoscience* 10, 366-369, doi:10.1038/ngeo2934 (2017).

444 43 Mouginot, J., Rignot, E., Scheuchl, B. & Millan, R. Comprehensive annual ice sheet velocity  
445 mapping using Landsat-8, Sentinel-1, and RADARSAT-2 data. *Remote Sens.* 9,  
446 doi:10.3390/rs9040364 (2017).

447 44 King, M. D. et al. Seasonal to decadal variability in ice discharge from the Greenland Ice  
448 Sheet. *Cryosphere* 12, 3813-3825, doi:10.5194/tc-12-3813-2018 (2018).

449 45 Porter, D. F. et al. Identifying Spatial Variability in Greenland's Outlet Glacier Response to  
450 Ocean Heat. *Front. Earth Sci.* 6, doi:10.3389/feart.2018.00090 (2018).

451 46 Rignot, E. & Mouginot, J. Ice flow in Greenland for the International Polar Year 2008-2009.  
452 *Geophysical Research Letters* 39, doi:10.1029/2012GL051634 (2012).

453 47 Sorensen, L. S. et al. Mass balance of the Greenland ice sheet (2003-2008) from ICESat data -  
454 the impact of interpolation, sampling and firn density. *Cryosphere* 5, 173-186 (2011).

455 48 Zwally, H. J. et al. Greenland ice sheet mass balance: distribution of increased mass loss with  
456 climate warming; 2003-07 versus 1992-2002. *Journal of Glaciology* 57, 88-102 (2011).

457 49 Rosenau, R., Scheinert, M. & Dietrich, R. A processing system to monitor Greenland outlet  
458 glacier velocity variations at decadal and seasonal time scales utilizing the Landsat imagery.  
459 *Remote Sensing of Environment* 169, 1-19, doi:10.1016/j.rse.2015.07.012 (2015).

460 50 Gogineni, S. et al. Coherent radar ice thickness measurements over the Greenland ice sheet.  
461 *Journal of Geophysical Research* 106, 33761-33772 (2001).

462 51 Shepherd, A. et al. Mass balance of the Antarctic Ice Sheet from 1992 to 2017. *Nature* 558,  
463 219-222, doi:10.1038/s41586-018-0179-y (2018).

464 52 Khan, S. A. et al. Geodetic measurements reveal similarities between post-Last Glacial  
465 Maximum and present-day mass loss from the Greenland ice sheet. *Sci. Adv.* 2,  
466 doi:10.1126/sciadv.1600931 (2016).

467 53 Peltier, W. R., Argus, D. F. & Drummond, R. Space geodesy constrains ice age terminal  
468 deglaciation: The global ICE-6G-C (VM5a) model. *J. Geophys. Res. B Solid Earth* 120, 450-487,  
469 doi:10.1002/2014JB011176 (2015).

470 54 Ettema, J. et al. Higher surface mass balance of the Greenland ice sheet revealed by high-  
471 resolution climate modeling. *Geophysical Research Letters* 36, doi:10.1029/2009GL038110  
472 (2009).

473 55 Bolch, T. et al. Mass loss of Greenland's glaciers and ice caps 2003-2008 revealed from  
474 ICESat laser altimetry data. *Geophysical Research Letters* 40, 875-881, doi:10.1002/grl.50270  
475 (2013).

476 56 Colgan, W., Luthcke, S., Abdalati, W. & Citterio, M. Constraining grace-derived cryosphere-  
477 attributed signal to irregularly shaped ice-covered areas. *Cryosphere* 7, 1901-1914,  
478 doi:10.5194/tc-7-1901-2013 (2013).

479 57 Tedesco, M. et al. Evidence and analysis of 2012 Greenland records from spaceborne  
480 observations, a regional climate model and reanalysis data. *The Cryosphere* 7, 615-630,  
481 doi:10.5194/tc-7-615-2013 (2013).

482 58 Vernon, C. L. et al. Surface mass balance model intercomparison for the Greenland ice sheet.  
483 *The Cryosphere* 7, 599-614, doi:10.5194/tc-7-599-2013 (2013).

484 59 Noël, B. et al. Modelling the climate and surface mass balance of polar ice sheets using  
485 RACMO2 - Part 1: Greenland (1958-2016). *Cryosphere* 12, 811-831, doi:10.5194/tc-12-811-  
486 2018 (2018).

487 60 Nghiem, S. V. et al. The extreme melt across the Greenland ice sheet in 2012. *Geophysical*  
488 *Research Letters* 39, doi:10.1029/2012GL053611 (2012).

489 61 Joughin, I., Abdalati, W. & Fahnestock, M. Large fluctuations in speed on Greenland's  
490 Jakobshavn Isbrae glacier. *Nature* 432, 608-610 (2004).

491 62 Joughin, I. et al. Continued evolution of Jakobshavn Isbrae following its rapid speedup. *J.*  
492 *Geophys. Res. F Earth Surf.* 113, doi:10.1029/2008JF001023 (2008).

493 63 Joughin, I., Smith, B. E. & Howat, I. Greenland Ice Mapping Project: Ice flow velocity variation  
494 at sub-monthly to decadal timescales. *Cryosphere* 12, 2211-2227, doi:10.5194/tc-12-2211-  
495 2018 (2018).

496 64 Rignot, E. Changes in ice dynamics and mass balance of the Antarctic ice sheet. *Philosophical  
497 Transactions of the Royal Society A-Mathematical Physical and Engineering Sciences* 364,  
498 1637-1655 (2006).

499 65 Howat, I. M., Joughin, I., Fahnestock, M., Smith, B. E. & Scambos, T. A. Synchronous retreat  
500 and acceleration of southeast Greenland outlet glaciers 2000-06: Ice dynamics and coupling  
501 to climate. *Journal of Glaciology* 54, 646-660 (2008).

502 66 Luckman, A., Murray, T., de Lange, R. & Hanna, E. Rapid and synchronous ice-dynamic  
503 changes in East Greenland. *Geophysical Research Letters* 33, art-L03503 (2006).

504 67 Dieng, H. B., Cazenave, A., Meyssignac, B. & Ablain, M. New estimate of the current rate of  
505 sea level rise from a sea level budget approach. *Geophysical Research Letters* 44, 3744-3751,  
506 doi:10.1002/2017GL073308 (2017).

507 68 Shepherd, A. & Nowicki, S. Improvements in ice-sheet sea-level projections. *Nat. Clim.  
508 Change* 7, 672-674, doi:10.1038/nclimate3400 (2017).

509 69 Fettweis, X. et al. Estimating the Greenland ice sheet surface mass balance contribution to  
510 future sea level rise using the regional atmospheric climate model MAR. *Cryosphere* 7, 469-  
511 489, doi:10.5194/tc-7-469-2013 (2013).

512 70 Markus, T. et al. The Ice, Cloud, and land Elevation Satellite-2 (ICESat-2): Science  
513 requirements, concept, and implementation. *Remote Sensing of Environment* 190, 260-273,  
514 doi:10.1016/j.rse.2016.12.029 (2017).

515 71 Flechtner, F. et al. What Can be Expected from the GRACE-FO Laser Ranging Interferometer  
516 for Earth Science Applications? *Surveys in Geophysics* 37, 453-470, doi:10.1007/s10712-015-  
517 9338-y (2016).

## 518 Methods

### 519 Data

520 In this assessment we analyse 5 groups of data: estimates of ice sheet mass-balance determined  
521 from 3 distinct classes of satellite observations - altimetry, gravimetry and the input–output method  
522 (IOM) - and model estimates of surface mass balance (SMB) and glacial isostatic adjustment (GIA).  
523 Each dataset is computed following previously reported methods (Supplementary Table 1) and, for  
524 consistency, they are aggregated within common spatial and temporal domains. Altogether, 26  
525 separate ice sheet mass balance datasets were used - 9 derived from satellite altimetry, 3 derived  
526 from the input-output method, and 14 derived from satellite gravimetry - with a combined period  
527 running from 1992 to 2018 (Extended Data Figure 1). We also assess 6 model estimates of GIA  
528 (Extended Data Table 1) and 10 model estimates of SMB (Extended Data Table 2).

### 529 Drainage Basins

530 We analyse mass trends using two ice sheet drainage basin sets (Extended Data Figure 2), to allow  
531 consistency with those used in the first IMBIE assessment <sup>1</sup>, and to evaluate an updated definition  
532 tailored towards mass budget assessments. The first set comprises 19 drainage basins delineated  
533 using surface elevation maps derived from ICESat-1 with a total area of 1,703,625 km<sup>2</sup> <sup>21</sup>. The second

drainage basin set is an updated definition considering other factors such as the direction of ice flow and includes 6 basins with a combined area of 1,723,300 km<sup>2</sup> <sup>46</sup>. The two drainage basin sets differ by 1% in area at the scale of the Greenland Ice Sheet, and this has a negligible impact on mass trends when compared to the estimated uncertainty of individual techniques.

### Glacial isostatic adjustment

GIA - the delayed response of Earth's interior to temporal changes in ice loading - affects estimates of ice sheet mass balance determined from satellite gravimetry and, to a lesser extent, satellite altimetry <sup>72</sup>. Here, we compare 6 independent models of GIA in the vicinity of the Greenland Ice Sheet (Extended Data Table 1). The GIA model solutions differ for a variety of reasons, including differences in their physics, in their computational approach, in their prescriptions of solid Earth unloading during the last glacial cycle and their Earth rheology, and in the data sets against which they are evaluated. No approach is generally accepted as optimal, and so we evaluate the models by computing the mean and standard deviation of their predicted uplift rates (Extended Data Figure 3). We also estimate the contribution of each model to gravimetric mass trends using a common processing approach <sup>51</sup> which puts special emphasis on the treatment of low spherical harmonic degrees in the GIA-related trends in the gravitational field.

The highest rates of GIA-related uplift occur in northern Greenland - though this region also exhibits marked variability among the solutions, as does the area around Kangerlussuaq Glacier to the southeast. Even though the model spread is high in northern Greenland, the signal in this sector is also consistently high in most solutions. However, none of the GIA models considered here fully captures all areas of high uplift present in the models, and so it is possible there is a bias towards low values in the average field across the ice sheet overall. The models yield an average adjustment for GRACE estimates of Greenland Ice Sheet mass balance of -3 Gt/yr, with a standard deviation of around 20 Gt/yr. The spread is likely in part due to differences in the way each model accounts for GIA in North America which is ongoing and impacts western Greenland, and so care must be taken when estimating mass balance at basin scale. Local misrepresentation of the solid Earth response can also have a relatively large impact stemming especially from lateral variations of solid-Earth properties <sup>52</sup>, and revisions of the current state of knowledge can be expected <sup>41</sup>.

### Surface mass balance

Here, ice-sheet SMB is defined as total precipitation minus sublimation, evaporation and meltwater runoff, i.e. the interaction of the atmosphere and the superficial snow and firn layers, for example through mass exchanges via precipitation, sublimation, and runoff, and through mass redistribution by snowdrift, melting, and refreezing. We compare 10 estimates of Greenland Ice Sheet SMB derived using a range of alternative approaches; 4 regional climate models (RCM's), 2 downscaled RCM's, a global reanalysis, 2 downscaled model reanalyses of climate data, and 1 gridded model of snow processes driven by climate model output (Extended Data Table 2).

Although SMB models of similar class tend to produce similar results, there are larger differences between classes – most notably the global reanalysis and the process model which lead to estimates of SMB that are significantly higher and lower than all other solutions, respectively. The regional climate model solutions agree well at the scale of individual drainage sectors, with the largest differences occurring in north-east Greenland (Extended Data Figure 4). The snow process model tends to underestimate SMB when compared to the other solutions we have considered in various sectors of the ice sheet, at times even yielding negative SMB, while the global reanalysis tends to overestimate it.

Across all models, the average SMB of the Greenland Ice Sheet between 1980 to 2012 is 351 Gt/yr and the standard deviation is 98 Gt/yr. However, the spread among the 8 RCM's and downscaled reanalyses is considerably smaller; these solutions lead to an average Greenland Ice Sheet SMB of 361 Gt/yr with a standard deviation of 40 Gt/yr over the same period. By comparison, the global reanalysis and process model lead to ice sheet wide estimates of SMB that are significantly larger (504 Gt/yr) and smaller (125 Gt/yr) than this range, respectively. Model resolution is an important factor when estimating SMB and its components, as respective contributions where only the spatial resolution differed yield regional differences. Additionally, the underlying model domains were identified as a source of discrepancy in the case of the Greenland Ice Sheet, as some products would allocate the ablation area outside the given mask.

### Individual estimates of ice sheet mass balance

To standardise our comparison and aggregation of the 26 individual satellite estimates of Greenland Ice Sheet mass balance, we applied a common approach to derive rates of mass change from cumulative mass trends<sup>51</sup>. Rates of mass change were computed over 36-month intervals centred on regularly spaced (monthly) epochs within each cumulative mass trend time series, oversampling the individual time series where necessary. At each epoch, rates of mass change were estimated by fitting a linear trend to data within the surrounding 36-month time window using a weighted least-squares approach, with each point weighted by its measurement error. The associated mass trend uncertainties were estimated as the **root sum square of the regression error and the measurement error**. Time series were truncated by half the moving-average window period at the start and end of their period. The emerging rates of mass change were then averaged over 12-month periods to reduce the impact of seasonal cycles.

**Gravimetry** We include 14 estimates of Greenland Ice Sheet ice sheet mass balance determined from GRACE satellite gravimetry which together span the period 2003 to 2016 (Extended Data Figure 1). 10 of the gravimetry solutions were computed using spherical harmonic solutions to the global gravity field and 4 were computed using spatially defined mass concentration units (Supplementary Table 1). **An unrestricted range** of alternative GIA corrections were used in the formation of the gravimetry mass balance solutions based on commonly-adopted model solutions and their variants<sup>41,53,73-78</sup> (Supplementary Table 1). There was some variation in the sampling of the individual gravimetry data sets, and their collective effective (weighted mean) temporal resolution is 0.08 years. Overall, there is good agreement between rates of Greenland Ice Sheet mass change derived from satellite gravimetry (Extended Data Figure 5); all solutions show the ice sheet to be in a state of negative mass balance throughout their survey periods, with mass loss peaking in 2012 and reducing thereafter. **During the period 2005 to 2015**, annual rates of mass change determined from satellite gravimetry differ by **86 Gt/yr on average**, and their average standard deviation is 26 Gt/yr (Extended Data Table 3).

**Altimetry** We include 9 estimates of Greenland Ice Sheet mass balance determined from satellite altimetry which together span the period 2004 to 2018 (Extended Data Figure 1). 3 of the solutions are derived from radar altimetry, 4 from laser altimetry, and 2 use a combination of both (Supplementary Table 1). The altimetry mass trends are also computed using a range of approaches, including crossovers, planar fits, and repeat track analyses. The laser altimetry mass trends are computed from ICESat-1 data as constant rates of mass change over their respective survey periods, while the radar altimetry mass trends are computed from EnviSat and/or CryoSat-2 data with a temporal resolution of between 1 and 72 months. In consequence, the altimetry solutions have an effective collective temporal resolution of 0.74 years. Mass changes are computed after making corrections for alternative sources of surface elevation change, including glacial isostatic and elastic



adjustment, and firn height changes (see Supplementary Table 1). Despite the range of input data and technical approaches, there is good overall agreement between rates of mass change determined from the various satellite altimetry solutions (Extended Data Figure 5). All altimetry solutions show the Greenland Ice Sheet to be in a state of negative mass balance throughout their survey periods, with mass loss peaking in 2012 and reducing thereafter. During the period 2005 to 2015, annual rates of mass change determined from satellite altimetry differ by 84 Gt/yr on average, and, their average standard deviation is 31 Gt/yr (Extended Data Table 3). The greatest variance lies among the 4 laser altimetry mass balance solutions which range from -248 to -128 Gt/yr between 2004 and 2010; aside from methodological differences, possible explanations for this high spread include the relatively short period over which the mass trends are determined, the poor temporal resolution of these data sets, and the rapid change in mass balance occurring during the period in question.

**Input-Output Method** We include 3 estimates of Greenland Ice Sheet mass balance determined from the input-output method which together span the period 1992 to 2015 (Extended Data Figure 1). Although there are relatively few data sets by comparison to the gravimetry and altimetry solutions, the input-output data provide information on the partitioning of the mass change (surface processes and/or ice dynamics) cover a significantly longer period and are therefore an important record of changes in Greenland Ice Sheet mass during the 1990's. The input-output method makes use of a wide range of satellite imagery for computing ice sheet discharge (output), and several alternative SMB model estimates of snow accumulation (input) and runoff (output) (see Supplementary Table 1). 2 of the input-output method datasets exhibit temporal variability across their survey periods, and 2 provide only constant rates of mass changes. Although these latter records are relatively short, they are an important marker with which variances among independent estimates can be evaluated. The collective effective (weighted mean) temporal resolution of the input-output method data is 0.14 years, although it should be noted that in earlier years the satellite ice discharge component of the data are relatively sparsely sampled in time (e.g. <sup>79</sup>). There is good overall agreement between rates of mass change determined from the input-output method solutions (Extended Data Figure 5). During the period 2005 to 2015, annual rates of mass change determined from the 4 input-output data sets differ by up to 47 Gt/yr on average, and their average standard deviation is 22 Gt/yr (Extended Data Table 3). These differences are comparable to the estimated uncertainty of the individual techniques and are also small relative to the estimated mass balance over the period in question. In addition to showing that the Greenland Ice Sheet was in a state of negative mass balance since 2000, with mass loss peaking in 2012 and reducing thereafter, the input-output method data show that the ice sheet was close to a state of balance prior to this period <sup>40</sup>.

### Aggregate estimate of ice sheet mass balance

To produce an aggregate estimate of Greenland Ice Sheet mass balance, we combine the 14 gravimetry, 9 altimetry, and 3 input-output method datasets to produce a single 26-year record spanning the period 1992 to 2018. First, we combine the gravimetry, altimetry, and the input-output method data separately into three time-series by forming an error-weighted average of individual rates of ice sheet mass change computed using the same technique (Extended Data Figure 6). At each epoch, we estimate the uncertainty of these time-series as the root mean square of their component time-series errors. We then combine the mass balance time-series derived from gravimetry, altimetry, and the input-output method to produce a single, aggregate (reconciled) estimate, computed as the error-weighted mean of mass trends sampled at each epoch. We estimated the uncertainty of this reconciled rate of mass balance as either the root mean square

departure of the constituent mass trends from their weighted-mean or the root mean square of their uncertainties, whichever is larger, divided by the square root of the number of independent satellite techniques used to form the aggregate. Cumulative uncertainties are computed as the root sum square of annual errors, on the assumption that annual errors are not correlated over time. This assumption has been employed in numerous mass balance studies<sup>1,18,40,51</sup>, and its effect is to reduce cumulative errors by a factor 2.2 over the 5-year periods we employ in this study (Table 1). If some sources of error are temporally correlated, the cumulative uncertainty may therefore be underestimated. In a recent study, for example, it is estimated that 30 % of the annual mass balance error is systematic (Shepherd et al., 2019), and in this instance the cumulative error may be 37 % larger. On the other hand, the estimated annual error on aggregate mass trends reported in this study (61 Gt/yr) are 70% larger than the spread of the independent estimates from which they are combined (36 Gt/yr)(Extended Data Table 3), which suggests the underlying errors may be overestimated by a similar degree. A more detailed analysis of the measurement and systematic errors is required to improve the cumulative error budget.

During the period 2004 to 2015, when all three satellite techniques were in operation, there is good agreement between changes in ice sheet mass balance on a variety of timescales (Extended Data Figure 6). In Greenland, there are large annual cycles in mass superimposed on equally prominent interannual fluctuations as well as variations of intermediate (~5 years) duration. These signals are consistent with fluctuations in SMB that have been identified in meteorological records<sup>1,80</sup>, and are present within the time-series of mass balance emerging from all three satellite techniques, to varying degrees, according to their effective temporal resolution. For example, correlated seasonal cycles are apparent in the gravimetry and input-output method mass balance time series, because their effective temporal resolutions are sufficiently short (0.08 and 0.14 years, respectively) to resolve such changes. However, at 0.74 years, the effective temporal resolution of the altimetry mass balance time series is too coarse to detect cycles on sub-annual timescales. Nevertheless, when the aggregated mass balance data emerging from all three experiment groups are degraded to a common temporal resolution of 36 months, the time-series are well correlated ( $0.63 < r^2 < 0.80$ ) and, over longer periods, all techniques identify the marked increases in Greenland Ice Sheet mass loss peaking in 2012. During the period 2005 to 2015, annual rates of mass change determined from all three techniques differ by up to 137 Gt/yr on average, and their average standard deviation is 36 Gt/yr - a value that is small or compared to their estimated uncertainty (61 Gt/yr)(Extended Data Table -3).

## Data availability

The aggregated Greenland Ice Sheet mass-balance data generated in this study are freely available at <http://www.imbie.org/data-downloads>.

## Extended Data Legends

**Extended Data Figure 1** | Ice sheet mass balance data sets used in this study and their main contributors (**top**) and the number and class of data available in each calendar year (**bottom**). The interval 2003 to 2010 includes almost all datasets and is selected as the overlap period. Further details of the satellite observations used in this study are provided in Supplementary Table 1.

**Extended Data Figure 2** | Greenland Ice Sheet drainage basins used in this study, according to the definitions of ref <sup>21</sup> (left) and ref <sup>46</sup> (right).

**Extended Data Figure 3** | Bedrock uplift rates in Greenland averaged over the glacial isostatic adjustment (GIA) model solutions used in this study (left), as well as their standard deviation (right). Further details of the GIA models used in this study are provided in Extended Data Table 1. High rates of uplift and subsidence associated with the former Laurentide Ice Sheet are apparent to the southwest of Greenland.

**Extended Data Figure 4** | Time series of surface mass balance (SMB) in Greenland Ice Sheet drainage basins <sup>81,82</sup>. Solid lines are annual averages of the monthly data (dashed lines). Further details of the SMB models used in this study are provided in Extended Data Table 2.

**Extended Data Figure 5** | Individual rates of Greenland ice-sheet mass balance used in this study as determined from satellite altimetry (a, left), gravimetry (b, centre) and the input–output method (c, right). The light-grey shading shows the estimated 1 $\sigma$  uncertainty relative to the ensemble average. The standard error of the mean solutions, per epoch, is shown in mid-grey.

**Extended Data Figure 6** | Rate of Greenland Ice Sheet mass balance as derived from the three techniques of satellite radar and laser altimetry (red), input-output method (blue), and gravimetry (green), and their arithmetic mean (gray), with uncertainty ranges (light shading).

**Extended Data Figure 7** | **Cumulative Greenland Ice Sheet surface mass balance.** The cumulative surface mass change (lightest blue) determined from an average of the RACMO2.3p2 <sup>59</sup> (light blue), MARv3.6 <sup>22</sup> (mid-blue) and HIRHAM <sup>9</sup> (dark blue) regional climate models relative to their 1980-1990 means (see Methods). The estimated uncertainty of the average change is also shown (shaded area) is computed as the average of the uncertainties from each of the three models. RACMO2.3p2 uncertainties are based upon a comparison to in-situ observations <sup>40</sup>. MARv3.6 uncertainties are evaluated from the variability due to forcing from climate reanalyses <sup>22</sup>. HIRHAM uncertainties are estimated based on comparisons to in-situ accumulation and ablation data <sup>83</sup>. Cumulative uncertainties are computed as the root sum square of annual errors, on the assumption that these errors are not correlated over time <sup>18</sup>.

**Extended Data Table 1.** Details of Glacial Isostatic Adjustment (GIA) models used in this study.

†Regional changes in mass associated with the GIA signal determined by the contributor.

‡Regional changes in mass associated with the GIA signal calculated as an indicative rate using spherical-harmonic degrees 3 to 90 and a common treatment of degree 2<sup>88</sup>.

<sup>a</sup> Main reference publication(s).

<sup>b</sup> Model from main publication unless otherwise stated. Comma-separated values refer to properties of a radially varying (1D, one-dimensional) Earth model: the first value is lithosphere thickness (km), other values reflect mantle viscosity ( $\times 10^{21}$  Pa s) for specific layers; see relevant publication.

<sup>c</sup> GIA model details: SH=spherical harmonic (maximum degree indicated), FE=finite element, C=compressible, IC=incompressible, RF=rotational feedback, SG=self-gravitation, OL=ocean loading, 'x' = feature not included.

<sup>d</sup> RSL = relative sea-level data; GPS rates corrected for elastic response to contemporary ice mass change.

<sup>e</sup> Earth model taken from ref<sup>53</sup>

<sup>f</sup> Ice model taken from ref<sup>53</sup>

<sup>g</sup> Different to ICE-6G\_C in Antarctica, owing to the use of BEDMAP2<sup>89</sup> topography.

**Extended Data Table 2.** Details of the surface mass balance (SMB) models used in this study.


<sup>a</sup> Main reference publication; additional references are provided in Supplementary Table 1.


<sup>b</sup> SMB model class; regional climate model (RCM), global numerical analysis (GA), process model (PM). Native resolution (n) and downscaled (d) models are also identified.

<sup>c</sup> Averages over the period 1980 to 2012 for the Greenland Ice Sheet excluding peripheral ice caps and using the drainage basins from ref<sup>46</sup>.

**Extended Data Table 3:** Estimates of ice-sheet mass balance from satellite altimetry, gravimetry the input–output method, and from all three groups during the period 2005 to 2015. Also shown are the average standard deviations (s.d.) and ranges of individual estimates within each group during the same period.


\*No altimetry data in 2010.







Extended Data Figure 2









Extended Data Figure 4







Extended Data Figure 7

Contributor	Model	Publication <sup>a</sup>	Earth model <sup>b</sup>	Ice model <sup>b</sup>	GIA model <sup>c</sup>	Constraint data <sup>d</sup>	GIA (Gt/yr)
A	A13	<sup>76</sup>	VM5a (1D) <sup>e</sup>	ICE-6G_C <sup>f</sup>	SH, C, RF, SG, OL	As for ICE-6G_C <sup>f</sup>	-9 <sup>‡</sup>
Lecavalier	Huy3	<sup>41</sup>	1D (120, 0.5, 2)	Huy3/ICE-5G	SH(256), IC, RF, SG, OL	RSL, ice extent, paleo thinning rates	-19 <sup>‡</sup>
Sasgen	GGG1D.0	<sup>52,84</sup>	VM-GPS <sup>52</sup>	modified GREEN1 <sup>85</sup>	SH(256)/FE(radial), IC, RF, SG, OL	GPS, RSL	+17 <sup>‡</sup>
Peltier	ICE-6G_D (VM5a)	<sup>53</sup>	VM5a (1D) <sup>e</sup>	ICE-6G_D <sup>g</sup>	SH(512)	GPS, RSL, Earth rotation	-10 <sup>‡</sup>
van der Wal	SL-dry-4mm/W12	<sup>86</sup>	3D, power-law rheology	Combination of W12 (Antarctica) and ICE-5G	FE, IC, xRF	GPS, RSL, seismic velocities (Earth model)	+21 <sup>‡</sup>
Spada	SELEN 4	<sup>87</sup>	VM5a (3-layer average of 1D model) <sup>e</sup>	ICE-6G_C <sup>f</sup>	SELEN4: SH(128), IC, RF, SG, OL	As for ICE-6G_C <sup>f</sup>	-27 <sup>‡</sup>
Extended Data Table 1							



Contributor	Model	Publication <sup>a</sup>	Class <sup>b</sup>	Area (10 <sup>6</sup> km <sup>2</sup> )	Grid	SMB <sup>c</sup> (Gt/yr)	Precipitation <sup>c</sup> (Gt/yr)	Runoff <sup>c</sup> (Gt/yr)
Noël	RACMO2.3	<sup>90</sup>	RCM	1.73	11 km	350	721	311
Noël	RACMO2.3p2	<sup>59</sup>	RCM	1.73	11 km	432	727	258
Langen	HIRHAM5	<sup>9</sup>	RCM	1.71	5.5 km	385	794	351
Fettweis	MARv3.6	<sup>22</sup>	RCM	1.69	7.5 km	381	706	308
Noël	RACMO2.3d	<sup>91</sup>	RCM-d	1.69	1 km	314	755	397
Noël	RACMO2.3p2d	<sup>59</sup>	RCM-d	1.69	1 km	338	703	331
Cullather	MERRA-2	<sup>92</sup>	GA-n	1.73	0.5 °	504	818	277
Hanna	ECMWF	<sup>14</sup>	GA-d	1.65	5 km	370	532	186
Wilton	ECMWFd	<sup>93</sup>	GA-d	1.71	1 km	314	603	246
Mernild	Snow Model	<sup>94</sup>	PM	1.64	5 km	125	655	418
Extended Data Table 2								

786

Technique	Mass balance (Grt/yr)	s.d.(Gt/yr)	range (Gt/yr)
Altimetry*	-237 ± 46	31	84
Gravimetry	-250 ± 52	26	86
Input-Output Method	-269 ± 80	22	47
All	-249 ± 61	36	137
Extended Data Table 3			

787

788    [Supplementary Information](#)

This table is an excel spreadsheet	
<b>Supplementary Table 1</b>	This table contains details of the satellite datasets used in this study <sup>35,36,40,73,77,80,95-118</sup> .

789

## Additional References

- 72 Wahr, J., Wingham, D. & Bentley, C. A method of combining ICESat and GRACE satellite data to constrain Antarctic mass balance. *Journal of Geophysical Research-Solid Earth* **105**, 16279-16294 (2000).
- 73 Paulson, A., Zhong, S. & Wahr, J. Inference of mantle viscosity from GRACE and relative sea level data. *Geophysical Journal International* **171**, 497-508, doi:10.1111/j.1365-246X.2007.03556.x (2007).
- 74 Peltier, W. R. in *Annual Review of Earth and Planetary Sciences* Vol. 32 111-149 (2004).
- 75 Simpson, M. J. R., Milne, G. A., Huybrechts, P. & Long, A. J. Calibrating a glaciological model of the Greenland ice sheet from the Last Glacial Maximum to present-day using field observations of relative sea level and ice extent. *Quaternary Science Reviews* **28**, 1631-1657, doi:10.1016/j.quascirev.2009.03.004 (2009).
- 76 A, G., Wahr, J. & Zhong, S. Computations of the viscoelastic response of a 3-D compressible earth to surface loading: An application to glacial isostatic adjustment in Antarctica and Canada. *Geophysical Journal International* **192**, 557-572, doi:10.1093/gji/ggs030 (2013).
- 77 Schrama, E. J. O., Wouters, B. & Rietbroek, R. A mascon approach to assess ice sheet and glacier mass balances and their uncertainties from GRACE data. *J. Geophys. Res. B Solid Earth* **119**, 6048-6066, doi:10.1002/2013JB010923 (2014).
- 78 Klemann, V. & Martinec, Z. Contribution of glacial-isostatic adjustment to the geocenter motion. *Tectonophysics* **511**, 99-108, doi:10.1016/j.tecto.2009.08.031 (2011).
- 79 Rignot, E. *et al.* Recent Antarctic ice mass loss from radar interferometry and regional climate modelling. *Nature Geoscience* **1**, 106-110, doi:10.1038/ngeo102 (2008).
- 80 Wouters, B., Bamber, J. L., Van Den Broeke, M. R., Lenaerts, J. T. M. & Sasgen, I. Limits in detecting acceleration of ice sheet mass loss due to climate variability. *Nature Geoscience* **6**, 613-616, doi:10.1038/ngeo1874 (2013).
- 81 Rignot, E., Mouginot, J. & Scheuchl, B. Ice Flow of the Antarctic Ice Sheet. *Science* **333**, 1427-1430, doi:10.1126/science.1208336 (2011).
- 82 Rignot, E., Mouginot, J. & Scheuchl, B. Antarctic grounding line mapping from differential satellite radar interferometry. *Geophysical Research Letters* **38**, doi:10.1029/2011GL047109 (2011).
- 83 Langen, P. L., Fausto, R. S., Vandecrux, B., Mottram, R. H. & Box, J. E. Liquid water flow and retention on the Greenland ice sheet in the regional climate model HIRHAM5: Local and large-scale impacts. *Front. Earth Sci.* **4**, doi:10.3389/feart.2016.00110 (2017).
- 84 Martinec, Z. Spectral-finite element approach to three-dimensional viscoelastic relaxation in a spherical earth. *Geophysical Journal International* **142**, 117-141, doi:10.1046/j.1365-246X.2000.00138.x (2000).
- 85 Fleming, K. & Lambeck, K. Constraints on the Greenland Ice Sheet since the Last Glacial Maximum from sea-level observations and glacial-rebound models. *Quaternary Science Reviews* **23**, 1053-1077, doi:10.1016/j.quascirev.2003.11.001 (2004).
- 86 King, M. A., Whitehouse, P. L. & van der Wal, W. Incomplete separability of Antarctic plate rotation from glacial isostatic adjustment deformation within geodetic observations. *Geophysical Journal International* **204**, 324-330, doi:10.1093/gji/ggv461 (2016).
- 87 SELEN v2.9.13 [software] (2018).
- 88 Martinec, Z. & Hagedoorn, J. The rotational feedback on linear-momentum balance in glacial isostatic adjustment. *Geophysical Journal International* **199**, 1823-1846, doi:10.1093/gji/ggu369 (2014).
- 89 Fretwell, P. *et al.* Bedmap2: Improved ice bed, surface and thickness datasets for Antarctica. *Cryosphere* **7**, 375-393, doi:10.5194/tc-7-375-2013 (2013).

839 90 Noël, B. *et al.* Evaluation of the updated regional climate model RACMO2.3: Summer  
840 snowfall impact on the Greenland Ice Sheet. *Cryosphere* **9**, 1831-1844, doi:10.5194/tc-9-  
841 1831-2015 (2015).

842 91 Noël, B. *et al.* A daily, 1 km resolution data set of downscaled Greenland ice sheet surface  
843 mass balance (1958-2015). *Cryosphere* **10**, 2361-2377, doi:10.5194/tc-10-2361-2016 (2016).

844 92 Gelaro, R. *et al.* The modern-era retrospective analysis for research and applications, version  
845 2 (MERRA-2). *Journal of Climate* **30**, 5419-5454, doi:10.1175/JCLI-D-16-0758.1 (2017).

846 93 Wilton, D. J. *et al.* High resolution (1 km) positive degree-day modelling of Greenland ice  
847 sheet surface mass balance, 1870-2012 using reanalysis data. *Journal of Glaciology* **63**, 176-  
848 193, doi:10.1017/jog.2016.133 (2017).

849 94 Mernild, S. H., Liston, G. E., Hiemstra, C. A. & Christensen, J. H. Greenland Ice Sheet Surface  
850 Mass-Balance Modeling in a 131-Yr Perspective, 1950-2080. *Journal of Hydrometeorology*  
851 **11**, 3-25 (2010).

852 95 Bonin, J. & Chambers, D. Uncertainty estimates of a GRACE inversion modelling technique  
853 over greenland using a simulation. *Geophysical Journal International* **194**, 212-229,  
854 doi:10.1093/gji/ggt091 (2013).

855 96 Blazquez, A. *et al.* Exploring the uncertainty in GRACE estimates of the mass redistributions  
856 at the Earth surface: Implications for the global water and sea level budgets. *Geophysical*  
857 *Journal International* **215**, 415-430, doi:10.1093/gji/ggy293 (2018).

858 97 Forsberg, R., Sørensen, L. & Simonsen, S. Greenland and Antarctica Ice Sheet Mass Changes  
859 and Effects on Global Sea Level. *Surveys in Geophysics* **38**, 89-104, doi:10.1007/s10712-016-  
860 9398-7 (2017).

861 98 Groh, A. & horwath, M. in *European Geophysical Union* (Vienna, 2016).

862 99 Harig, C. & Simons, F. J. Mapping Greenland's mass loss in space and time. *Proceedings of*  
863 *the National Academy of Sciences of the United States of America* **109**, 19934-19937,  
864 doi:10.1073/pnas.1206785109 (2012).

865 100 Luthcke, S. B. *et al.* Antarctica, Greenland and Gulf of Alaska land-ice evolution from an  
866 iterated GRACE global mascon solution. *Journal of Glaciology* **59**, 613-631,  
867 doi:10.3189/2013JoG12J147 (2013).

868 101 Andrews, S. B., Moore, P. & King, M. A. Mass change from GRACE: A simulated comparison  
869 of Level-1B analysis techniques. *Geophysical Journal International* **200**, 503-518,  
870 doi:10.1093/gji/ggu402 (2015).

871 102 Save, H., Bettadpur, S. & Tapley, B. D. High-resolution CSR GRACE RL05 mascons. *J. Geophys.*  
872 *Res. B Solid Earth* **121**, 7547-7569, doi:10.1002/2016JB013007 (2016).

873 103 Seo, K. W. *et al.* Surface mass balance contributions to acceleration of Antarctic ice mass loss  
874 during 2003-2013. *J. Geophys. Res. B Solid Earth* **120**, 3617-3627, doi:10.1002/2014JB011755  
875 (2015).

876 104 Velicogna, I., Sutterley, T. C. & Van Den Broeke, M. R. Regional acceleration in ice mass loss  
877 from Greenland and Antarctica using GRACE time-variable gravity data. *Geophysical*  
878 *Research Letters* **41**, 8130-8137, doi:10.1002/2014GL061052 (2014).

879 105 Vishwakarma, B. D., Horwath, M., Devaraju, B., Groh, A. & Sneeuw, N. A Data-Driven  
880 Approach for Repairing the Hydrological Catchment Signal Damage Due to Filtering of GRACE  
881 Products. *Water Resources Research* **53**, 9824-9844, doi:10.1002/2017WR021150 (2017).

882 106 Wiese, D. N., Landerer, F. W. & Watkins, M. M. Quantifying and reducing leakage errors in  
883 the JPL RL05M GRACE mascon solution. *Water Resources Research* **52**, 7490-7502,  
884 doi:10.1002/2016WR019344 (2016).

885 107 Cheng, M., Tapley, B. D. & Ries, J. C. Deceleration in the Earth's oblateness. *J. Geophys. Res.*  
886 *B Solid Earth* **118**, 740-747, doi:10.1002/jgrb.50058 (2013).

887 108 Swenson, S., Chambers, D. & Wahr, J. Estimating geocenter variations from a combination of  
888 GRACE and ocean model output. *J. Geophys. Res. B Solid Earth* **113**,  
889 doi:10.1029/2007JB005338 (2008).

- 109 Csatho, B. M. *et al.* Laser altimetry reveals complex pattern of Greenland Ice Sheet dynamics. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 18478-18483, doi:10.1073/pnas.1411680112 (2014).
- 110 Nilsson, J., Gardner, A., Sørensen, L. S. & Forsberg, R. Improved retrieval of land ice topography from CryoSat-2 data and its impact for volume-change estimation of the Greenland Ice Sheet. *Cryosphere* **10**, 2953-2969, doi:10.5194/tc-10-2953-2016 (2016).
- 111 Gourmelen, N. *et al.* CryoSat-2 swath interferometric altimetry for mapping ice elevation and elevation change. *Advances in Space Research* **62**, 1226-1242, doi:10.1016/j.asr.2017.11.014 (2018).
- 112 Gunter, B. C. *et al.* Empirical estimation of present-day Antarctic glacial isostatic adjustment and ice mass change. *Cryosphere* **8**, 743-760, doi:10.5194/tc-8-743-2014 (2014).
- 113 Felikson, D. *et al.* Comparison of Elevation Change Detection Methods from ICESat Altimetry over the Greenland Ice Sheet. *IEEE Transactions on Geoscience and Remote Sensing* **55**, 5494-5505, doi:10.1109/TGRS.2017.2709303 (2017).
- 114 Helm, V., Humbert, A. & Miller, H. Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2. *Cryosphere* **8**, 1539-1559, doi:10.5194/tc-8-1539-2014 (2014).
- 115 Kjeldsen, K. K. *et al.* Improved ice loss estimate of the northwestern Greenland ice sheet. *J. Geophys. Res. B Solid Earth* **118**, 698-708, doi:10.1029/2012JB009684 (2013).
- 116 Khan, S. A. *et al.* Sustained mass loss of the northeast Greenland ice sheet triggered by regional warming. *Nat. Clim. Change* **4**, 292-299, doi:10.1038/nclimate2161 (2014).
- 117 Andersen, M. L. *et al.* Basin-scale partitioning of Greenland ice sheet mass balance components (2007-2011). *Earth and Planetary Science Letters* **409**, 89-95, doi:10.1016/j.epsl.2014.10.015 (2015).
- 118 Colgan, W. *et al.* Greenland ice sheet mass balance assessed by PROMICE (1995–2015). *GEUS Bulletin* **43** (2019).